

## **REVIEW**

# **Critique of medicinal conspicuousness of Parsley (*Petroselinum crispum*): A culinary herb of Mediterranean region**

**Sidra Mahmood<sup>1</sup>, Shahzad Hussain\*<sup>2</sup> and Farnaz Malik<sup>2</sup>**

<sup>1</sup>Department of Bioinformatics and Biotechnology, International Islamic University, Islamabad, Pakistan

<sup>2</sup>Drugs Control and Traditional Medicines Division, National Institute of Health, Islamabad, Pakistan

**Abstract:** WHO estimates, around 80% of the especially developing world is indigent on complementary and alternative medicines which are prodigiously derived from herbal material. Parsley (*Petroselinum crispum*) is an important culinary herb originated from the Mediterranean region. It possesses small and dark seeds with volatile oil content. *Petroselinum crispum* is now planted throughout the world due to its usage in food industry, perfume manufacturing, soaps, and creams. Its main constituents subsume coumarins, furanocoumarins (bergapten, imperatori), ascorbic acid, carotenoids, flavonoids, apiole, various terpenoid compounds, phenyl propanoids, phthalides, and tocopherol. Due to these constituents, it has been announced to possess a number of possible medicinal emblematics including, antimicrobial, antianemic, menorrhagic, anticoagulant, antihyperlipidemic, antihepatotoxic, antihypertensive, diuretic effects, hypoglycaemic, hypouricemic, anti oxidative and estrogenic activities. In Morocco, Parsley is mostly used as an elixir to treat arterial hypertension, diabetes, cardiac and renal diseases. Antioxidant and antibacterial activities of parsley, made it propitious in food systems. Its ELI17 gene has been corroborated as a particularly fast-responding gene. There is a requisite for extensive research to avail the maximal benefits of this significant medicinal plant. The aim of this review paper is to divulge the chemical constituents of parsley that are explicitly related to substantial medicinal facets.

**Keywords** *Petroselinum crispum*, hypoglycaemic, aggregation, apigenin.

## **INTRODUCTION**

Interest in the herbs has been escalating day by day, as they might offer a natural armor against the augmentation of certain maladies and be a conjectural treatment for some diseases. Parsley (*Petroselinum crispum*) is an important culinary herb commenced from the banks of Mediterranean basin, where it still can be found in wild forms. Parsley belongs to the genus *Petroselinum* Hill of the Apiaceae ancestors with the intention of being manipulated in the foodstuff, pharmaceutical, perfume and cosmetic productions (Lopez *et al.*, 1999). In Turkey, it is extensively disected and sprouted inside gardens and fields. New, dehydrated, and dried out foliage are utilized as a condiment, embellish and flavoring ingredient (Petropoulos *et al.*, 2004). Parsley (*Petroselinum crispum*) possesses small and dark seeds with volatile oil content (Tahan and Bayram, 2011). The indispensable oil of parsley has been reported to be immoderately manipulated in the victuals production and still as a cologne in perfume manufacturing, soaps, and creams nevertheless the oil for such functions is garnered from the seeds for the reason that the leaves yield minor amount. Parsley is propagated in the United States once a year, for its scented and beautiful foliage

(Simon and Quinn, 1988). Three major categories of parsley have been studied: 1) Unadorned leaf category (ssp neapolitanum, Danert), 2) Twisted leaf form (ssp crispum), primarily grown intended for their plants, 3) turnip- deep-rooted or 'Hamburg' category (ssp. tuberosum), which is cultivated designed for its pedigree (Petropoulos *et al.*, 2004). For the reason that of the soaring water substance (75-80%), parsley is customarily dehydrated for make known, in order to impede germs escalation and prevent deprivation due to biochemical feedback (Diaz *et al.*, 2002). It has been noted that aeration procedure of parsley foliage from *Petroselinum crispum* L. can sway the sensual and scented attributes of this herbal artifact. HPLC analysis actuated the 6"-O-malonylapiin to apiin fraction to be a appropriate indicator method. This fraction is altitudinous for unmarked and lyophilised foliage substance; oven-ventilation leads to demalonylation and consequently, to a squatty malonylapiin-apiin fraction (Lechtenberg *et al.*, 2007). Parsley plants and cell cultures are well recognized systems for studying non-host plant/pathogen interactions, indubitably with view to the titivation of phenyl propanoid metabolism (Hahlbrock and Scheel, 1989). The wastewater irrigation in parsley production fruitages socioeconomic benefits, but it also has negative effects on plant and musters heavy metal accumulation (Keser and Buyuk, 2012).

\*Corresponding author: e-mail: shshaikh2001@yahoo.com

### Chemical constituents


The parsley is found to have the following chemical constituents viz ascorbic acid, carotenoids, flavonoids, coumarins, apiole, various terpenoid compounds, phenyl propanoids, phthalides, furanocoumarins (bergapten, imperatori) and tocopherol have been chemically scouted (Tunali *et al.*, 1999). It has been reported that volatile compounds in unmarked parsley and in groups dried out, by means of the diverse drying techniques are following:  $\alpha$ -Pinene  $\beta$ -Myrcene,  $\alpha$ -Phellandrene,  $\beta$ -Phellandrene, cis-Ocimene, Isopropenyl-4-methylbenzene,  $\alpha$ -Terpinolene, p-Mentha-1,3,8-triene,  $\alpha$ -Copaene, Caryophyllene,  $\beta$ -Farnesen,  $\beta$ -Selinene,  $\gamma$ -Cadinene, Myristicin,  $\beta$ -Bisabolene,  $\beta$ -Sesquiphellandrene. The volatile oil of parsley seeds contain glycoside called apiin (apigenin-7-O-apiosyl-(1->2)-O-glucoside) (Tahan and Bayram, 2011).



Parsley (Damien *et al.*, 2011)




Seed capsules of *Petroselinum crispum* (blogspot.com) and Flowers of *Petroselinum crispum* (aphotoflora.com)



(Gregory *et al.*, 2012)

Cardinal oil constituents in dried parsley have been enunciated. These are as follows:  $\alpha$ -Thujene,  $\alpha$ -Pinene, Sabinene,  $\beta$ -Pinene,  $\beta$ -Myrcene,  $\alpha$ -Phellandrene, p-Cymene, Limonene,  $\beta$ -Phellandrene, (E)- $\beta$ -Ocimene,  $\gamma$ -Terpinene,  $\alpha$ -p-Dimethyl-phencone, p-cymenene, Phencone, Terpinolene, p-1,3,8-Menthatriene, Linalool, (Z) p-menth-2-en-1-ol, Dill ether, (Z) Dihydrocarvone, (E) Dihydrocarvone, (E) Carveol,  $\alpha$ -Terpineol, Estragol, Carvone, Bornyl acetate, (E,E) Decadienal,  $\alpha$ -Copaene, 2,5-Dimethyl-p-cymene, 2,5-Dimethoxy-p-


cymene, (E)  $\beta$ -Caryophyllene,  $\beta$ -Ionone, (Z) Anethole, Epi-bicyclosesquiphellandrene, Germacrene  $\alpha$ -Bergaptene, Myristicin,  $\alpha$ -Cadinol,  $\beta$ -Selinene,  $\alpha$ -Muurolene, Elemicin, Germacrene,  $\delta$ -Cadinol, Apiole, Phthalide isomer (vokk *et al.*, 2011). However, the most important oil components are p-mentha-1,3,8-triene, myristicin, apiole,  $\beta$ -phellandrene, myrcene, and isopropenyl-4-methylbenzene (Kasting *et al.*, 1972; Vernon and Richard 1983; MacLeod *et al.*, 1985).



Apiole

(Hui Zhang *et al.*, 2006)

Odorants of parsley leaves have been enumerated and mainly comprised of p-Mentha-1,3,8-triene, Myrcene, 2-Isopropyl-3-methoxypyrazine, 2-sec-Butyl-3-methoxypyrazine, Myristicin, Oct-1-en-3-one, (Z)-Octa-1,5-dien-3-one, Linalol, (E,E)-Deca-2,4-dienal, (Z)-Dec-6-enal, Methanethiol, (Z)-Hex-3-enal, p-Methylacetophenone, (Z)-Hex-3-enyl acetate, (Z)-Hex-3-enol, b-Phellandrene, 1-Isopropenyl-4-methylbenzene (Charly and Werner, 1998). Parsley aroma is substantially due to the domination of constituents viz. p-mentha-1,3,8-triene and myristicin (Jung *et al.*, 1992) in relationship with supplementary constituents already at extremely stumpy concentrations, specifically, 2-sec-butyl-3-methoxy-pyrazine, linalool, (Z) 6-decenal, and (Z) 3-hexenal (Masanetz and Grosch, 1998).




Myristicin

(Hui Zhang *et al.*, 2006)

*Petroselinum crispum* embodies ample amounts of the flavone apigenin (Justesen *et al.*, 1998) and the low concentration of other flavonoids. Flavones are a class of flavonoids discerned in a variety of fruits and vegetables and are most lavish in kumquats, parsley, and celery (Azzini *et al.*, 2007; Sakakibara *et al.*, 2003). Apigenin is a flavone espied in vegetables, seasonings (Ku' hnuu,


1976) and oranges (Fernandez de Simon *et al.*, 1992), and it heralds antioxidant activity *in vitro* (Fraga *et al.*, 1987; Van Acker *et al.*, 1996). Its potent biological effects have been elucidated *in vitro* and *in vivo*. Apigenin is imbibed from parsley, and exuded in low amounts, 0.58% on average, with urine. The parsley intervention resulted in sententious augmentation in the two antioxidant enzymes, GR (glutathione reductase) and SOD (superoxide dismutase) (Nielsen *et al.*, 1999).




R = H : Apigenin  
(Gregory *et al.*, 2012)

HPLC scrutiny of methanol extracts commencing elicitor-treated parsley cells, their manifestation through spectroscopic methods (MS, NMR) brandish two indication furanocoumarins marmesin and bergapten along with four structurally related phthalides 5-hydroxy-3-butylidenephthalide, 7-hydroxy-3-butylidenephthalide, 7-hydroxy-3-butylidenephthalide-7-O-glucoside and 7-hydroxy-3-butylidenephthalide-7-O-(6-malonylglucoside). Furanocoumarins are the foremost dispersible components contemplated to be powerful phytoalexins (Scheel *et al.*, 1986; Matern, 1991).

#### Furanocoumarins




Marmesin




Bergapten


#### Butylidenephthalides




5-hydroxybutylidenephthalide



7-Hydroxybutylidenephthalide




7-Hydroxybutylidenephthalide  
7-O-glucoside



7-Hydroxybutylidenephthalide  
7-O-(6-malonylglucoside)

(Jens *et al.*, 1999)

Through bioassay-guided separation, two benzoxoles, myristicin and apiole (Yakushijin *et al.*, 1983), two furocoumarins, cnidilin (Kuo-Hsiung *et al.*, 1969) and isoimperatorin, (Kuo-Hsiung *et al.*, 1969) and four flavone glycosides, apigetrin, (Redaelli *et al.*, 1980) apiin (Markham *et al.*, 1978) diosmetin 7-O-b-D-glucopyranoside and kaempferol 3-O-b-D-glucopyranoside (Okuyama *et al.*, 1978) together with a new glycoside, 6''-acetylapiin have been sequestered from American dried parsley. From Japanese fresh parsley, a new monoterpene glucoside called petroside has been quarantined together with apiin and icariside F<sub>2</sub>.



#### Pharmacological significance

Concerning 80% of the globe's populace is dependent on the exercise of conventional medication, which is commodiously supported on plant material (Dahiru *et al.*, 2006). Studies on a numeral of medicinal plants connote to facilitate propitious phytochemicals which know how to be mellowed for numerous fitness troubles (Gupta, 1994). In recent years, herbal medications encompass procured significance for the reason that of their standardization, effectiveness and charge value. These drugs are invariably single plant extracts or fractions thereof or mixtures of fractions and extracts from diverse plants (Subramoniam and Pushpangadan, 1991). Petroselinum crispum is now cultivated throughout the world and has been wielded in folk medicine as a diuretic, a stomachic, an emmenagogue and an abortifacient (Kreydiyyeh *et al.*, 2001).

Parsley has been promulgated to possess a number of possible medicinal idiosyncrasies including; antimicrobial (Wong and Kitts, 2006), antianemic, menorrhagic (Baytop, 1984), anticoagulant, antihyperlipidemic, antihepatotoxic, antihypertensive effects (Ozturk *et al.*, 1991), antioxidant (Nielsen *et al.*, 1999) as well as

laxative (Kreydiyyeh *et al.*, 2001). It has been bestowed to medicament lumbago, eczema, knee, ache, impotence, nose bleed and as a blood pressure regulator (Manderfeld *et al.*, 1997).

The seeds of parsley are also evinced like a diuretic owing to its elevated essential oil substance (Marczal *et al.*, 1997; Darias *et al.*, 2001). Parsley has a diuretic effect due to its constituent's apiol and myristicin (Newall *et al.*, 1994; Tyler, 1996; Kreydiyyeh and Usta, 2002) and because diuretics exaggerate the flow of urine, this is efficacious for the body to wash out bacteria as well as stones. Parsley accommodates calcium, iron, carotenes, ascorbic acid, and vitamin A (Louis, 1991). Because ascorbic acid is intimately renewed to oxalate, and furthermore emerge to augment the assimilation of nutritional oxalate, complementation may sprout the hazard of kidney pebbles (Massey *et al.*, 2005). Parsley leaf tea has no significant change on urine constitution or risk of urinary area rock creation in healthy subjects as well as there is a prerequisite for further research to reckon the consequences of parsley folio tea on urinary factors in fit and stone-developing patients (Fahad and Danny, 2011).

Hypoglycaemic activity of parsley has been unveiled by Ozsoy *et al.*, (2006). Its itemized antidiabetic functions may perhaps due to terpenoids (Pino *et al.*, 1997), flavonoid glucosides, coumarins (Anand *et al.*, 1981) otherwise ascorbic acid (Davey *et al.*, 1996). As an immemorial drug for diabetes, plant has been utilized in Turkey and in the whole world (Noel *et al.*, 1997). In folk medicine, it has been utilized to manipulate a commodious assortment of stipulations (Yanardag *et al.*, 2003a; Yanardag *et al.*, 2003a). It has also been conjectured, that parsley has antimicrobial activity and concentrate of parsley parade a noteworthy shielding outcome on the liver damage of diabetic rats (BolKent *et al.*, 2004; Manderfeld *et al.*, 1997). Antioxidant capacity has been symbolically aggrandized by subsidiary of the regime by means of unsullied parsley leaf can in rat plasma (Hempel *et al.*, 1999) and diminish oxidative hassle in humans (Nielsen *et al.*, 1999). Treatment of D-galactose-stressed mice with the ethanolic extract of *P. crispum* showed bulwark against the induced oxidative stress in brain regions. Therefore, it has been enunciated that parsley manifests a protective effect against mitochondrial oxidative damage in the mouse brain (Vora *et al.*, 2009). Actually, parsley contains voluminous hunk of flavonoids (apiin, luteolin, apigenin- glycosides), ascorbic acid, tocopherol and indispensable oils (apiole, myristicin) with anti-oxidant activities and these might preclude oxidative harm ( Marczal *et al.*, 1997; Nielsen *et al.*, 1999; Fejes *et al.*, 2000).


Apigenin has been imputed to be endowed with anti-carcinogenic (Wei *et al.* 1989; Birt *et al.* 1997), anti-inflammatory (Lee *et al.*, 1993) and anti-mutagenic

dominions (Kuo *et al.*, 1992). *In vitro* studies flaunt that the flavone apigenin of parsley restrains human lung, colon, breast, prostate, brain, and skin cancer cells; (Manthey *et al.*, 2002; Engelmann *et al.*, 2002) tongue cancer; (Walle *et al.*, 2007) and leukemia (Vargo *et al.*, 2006). Apigenin also abridge monocyte adhesion to LDL in vitro, showing potential to thwart one of the initial stages of atherosclerosis (Jeong *et al.*, 2007). In addition, animal studies with flavones substantiate the ability to alleviate the inflammatory response (Ueda *et al.*, 2004; Nicholas *et al.*, 2007).

It has been implied so as to parsley indispensable oil might be capable to restrain the cellular and humoral invulnerable retort. It knows how to decimate, evenly NO making as well as the purposes of macrophages as the central instinctive immune cells. Allergy, autoimmune along with chronic inflammatory disorders has been traditionally cured by parsley (Alireza *et al.*, 2012).

Intestinal disorders are ameliorated by parsley in the traditional herbal medicine. Parsley seed extract has inhibitory effect on ileum contraction and it was probably due to blocking of voltage- gated calcium channels (Moazedi *et al.*, 2007). It has been consummated that the aqueous and ethanol extracts of parsley endeavor antispasmodic activity on rat ileum. The relaxant effect of ethanolic extract was found better as compared to aqueous extract of parsley (Branković *et al.*, 2010) and non-selective antagonists of adrenoceptors (phentolamine and propranolol) do not stymie relaxation induced by extract. This justifies its use in folk medicine as a remedy for intestinal cramps and diarrhea (Irzaie *et al.*, 2010).

Parsley (*Petroselinum crispum*) and its major flavonol constituents Kaempferol and Quercetin are proficient of abating the uric acid levels in hyperuricemic. Therefore, the use of suboptimal dosages of allopurinol in combination with parsley intake may provide a snug avenue for prevention and treatment of hyperuricemia (Seid *et al.*, 2011).



Phenylketonuria (PKU) is an autosomal dormant genetic illness delineating through peculiarly lofty stages of phenylalanine in the transmission, along with the

nonappearance or noticeably curtailed enzymic properties of phenylalanine hydroxylase (PAH) in the liver (Folling, 1994; Jervis, 1960). It is induced by imperfections in the phenylalanine hydroxylase (PAH) genetic material (Folling, 1994). It has been used as an effectual therapeutic enzyme for the conduct of PKU (Ambrus *et al.*, 1978). It switches phenylalanine to harmless t-cinnamic acid (Hoskins *et al.*, 1984). The predicaments of this alluring enzyme treatment encompasses the little steadiness in the passage and the antigenicity of the plant enzyme. Recombinant PAL instigated from parsley (*Petroselinum crispum*) chemically conjugated with stimulated PEG2 [2, 4-bis (O-methoxypolyethy-leneglycol)-6-chloro-s-triazine] have been shown enormously augmented equanimity in the spread and abridges the plasma concentration of phenylalanine in mice. Therefore, PEG-PAL may be optimized for the protected acceptance stimulation by each element or genetic amendment of PAL, and can be bequeathed carefully to PKU patients devoid of the threat of destructive allergic reactions (Ikeda *et al.*, 2005).


In Morocco, parsley is frequently used to treat arterial hypertension (Ziyyat *et al.*, 1997; Eddouks *et al.*, 2002), diabetes, cardiac (Jouad *et al.*, 2001; Eddouks *et al.*, 2002) and renal diseases (Jouad *et al.*, 2001). Polyphenolic compounds,  $\alpha$  tocopherol and flavonoids of parsley could intercept platelet functions, activation and aggregation (Janssen *et al.*, 1998; Mabile *et al.*, 1999; Rein *et al.*, 2000; Pearson *et al.*, 2002;) furthermore fortress adjacent to cardiovascular maladies (Pace-Asciak *et al.*, 1995). Aqueous extract (Pc) of *Petroselinum crispum* (Mill), apigenin and cosmoiin possess a strong anti-platelet aggregation activity that interfere on haemostasis (Chaves *et al.*, 2011) causing an addendum of tail bleeding time and normalizing the platelet hyper-aggregability connected to cardiovascular disorders (Dounia *et al.*, 2009). Parsley also hindered *in vitro* thrombin and ADP-persuaded platelet gathering (Mekhfi *et al.*, 2004).

Insufficiency of internal estrogen (estrone, estradiol) secretion engenders several physical snarls shown in postmenopausal women, such as osteoporosis, blood cholesterol elevation, and symptoms of menopause (hot flashes and depression) (Masayuki *et al.*, 2000). The extracts of the aerial part of *Petroselinum crispum* MILL. (Parsley) that was cultivated in U.S.A. and Japan evinced reproducible estrogenic activity comparable to that of the isoflavone glucoside fraction (Kitagawa *et al.*, 1976) from soybean.

#### Food eminence upgrading

Parsley (*Petroselinum crispum*), is frequently employed to flavor the foods of China, Mexico, South America, India and South East Asia. Food quality deterioration majorly occurs due to lipid oxidation. Constituents of unsullied parsley folio exploration superoxide anion in

vitro (Campanella *et al.*, 2003) along with methanol concentrates of parsley scavenge hydroxyl fundamental in totaling to bulwarking in opposition to ascorbic acid-persuaded membrane oxidation (Fejes *et al.*, 2000). Phenolic composites hauled out from parsley have been found accountable, in fraction, for mutually antioxidant and antibacterial properties (Peter and David, 2006). The antioxidant activity of parsley in food systems is related to its total phenolic content and radical scavenging capacity but not to their propensity to chelate iron *in vitro* (Jimenez *et al.*, 2008).



Effects of Methanolic Extracts of Parsley, Apiin and Apigenin on uterus weight. It decreases in Ovariectomized Mice. Each column represents mean with the S.E. of 5 to 7 mice. (Masayuki *et al.*, 2000)

Parsley grown in summer time, could assist in mummifying foods because of colossal content of compounds with antimicrobial activity and a tenacious impeding effect of essential oils although different nutrient content of food may imprint microbial resistance (Díaz *et al.*, 2002). It has been noted for its antibacterial action in opposition to normal microflora, coliforms, yeast and molds and S.aureus in Kareish cheese, and its totaling is pertinent to the purchaser and might legacy to the improvement of novel and protected multifariousness of Kareish cheese (Wahba *et al.*, 2010).

#### Additional remarkable features

Parsley (*Petroselinum crispum*) ELI17 gene has been ascribed as a predominantly quick-responding gene emanating a new-fangled category of W box-containing, elicitor-responsive advertiser constituent, E17. The protein instructed by the ELI17 gene, demonstrates a variety of fundamental features of established transcription factors and is designated as a CMPG protein according to the first four strictly conserved amino acids, delineating a recently promising category of plant specific proteins (Christoph *et al.*, 2001). Phe ammonia-lyase (PAL) is an enzyme, which has been derived from parsley (*Petroselinum crispum*). For the reason that, its chief responsibility in secondary phenylpropanoid metabolism, it is solitary the main irregularly investigated plant enzymes (Holger and Georg, 2004). PAL actuate the impulsive, non-reductive amination of

**Table 1:** Imputing the discernible bioactive constituents along with their medicinal importance isolated from *Petroselinum crispum*.

| Chemical constituents  | Medicinal importance   | References   |
|--|--|--|
| Phenolic Compounds   | Antioxidant and antibacterial activities   | Peter and David, 2006.   |
| Apiol and Myristicin   | Diuretic effect  | Tyler, 1996; Newall <i>et al.</i> , 1994; Kreydiyyeh and Usta, 2002.   |
| Essential Oil Content  | Diuretic effect  | Darias <i>et al.</i> , 2001; Marczał <i>et al.</i> , 1997  |
| Ascorbic acid  | sprout the risk of kidney stones   | Massey <i>et al.</i> , 2005.   |
| Terpenoids<br>Flavonoid<br>Glucosides<br>Coumarins<br>Ascorbic acid  | Anti-diabetic properties   | Anand <i>et al.</i> , 1981; Davey <i>et al.</i> , 1996   |
| Flavonoids (apiin, luteolin, apigenin-glycosides), ascorbic acid, tocopherol and essential oils (apiole, myristicin) | Antioxidant properties   | Nielsen <i>et al.</i> , 1999; Fejes <i>et al.</i> , 2000; Marczał <i>et al.</i> , 1997.  |
| Flavone apigenin   | Restrains human lung, colon, breast, prostate, brain, and skin cancer cells, tongue cancer, Leukemia, initial stages of atherosclerosis. | Manthey <i>et al.</i> , 2002; Engelmann <i>et al.</i> , 2002; Walle <i>et al.</i> , 2007; Vargo <i>et al.</i> , 2006; Jeong <i>et al.</i> , 2007.        |
| Parsley essential oil  | Suppress the cellular and humoral immune response, allergy, autoimmune and chronic inflammatory disorders                                | Alireza <i>et al.</i> , 2012.  |
| Kaempferol and Quercetin   | Treatment of hyperuricemia   | Seid <i>et al.</i> , 2011.   |
| $\alpha$ tocopherol and flavonoids   | Intercept platelet functions, activation and aggregation, Cardiovascular diseases  | Mabile <i>et al.</i> , 1999; Janssen <i>et al.</i> , 1998 and Pearson <i>et al.</i> , 2002; Rein <i>et al.</i> , 2000; Pace-Asciak <i>et al.</i> , 1995. |
| Apigenin and cosmosiin   | Strong antiplatelet aggregation activity   | Chaves <i>et al.</i> , 2011.   |

trans-cinnamic acid to L-phenylalanine in the charisma of elevated ammonia congregations. By means of this asymmetric production, theoretical capitulates of 100% can be acquired; it is an enthralling response for manufacturing procedures (Sebastian and Uwe, 2010). Parsley seed oil has been found to endorse maximum inhibitory action in the direction of malonaldehyde (MA) configuration from squalene upon UV-irradiation (Wei and Shibamoto, 2007). Parsley sects distinguish the fungal *Phytophthora sojae*, a phytopathogen through a plasma covering receptor. This parsley receptor binds to a 13 amino acid oligopeptide portion (Pep-13) of a 42 k Da fungal cell barrier glycoprotein and instigates a multifaceted resistance comeback in cultured parsley cells (Dirk *et al.*, 1998).

## CONCLUSION

Parsley exhibits antimicrobial, antianemic, menorrhagic, anticoagulant, antihyperlipidemic, antihepatotoxic, antihypertensive ramifications, hypoglycemic, anti oxidative, estrogenic and hyperuricemic activity of parsley has been promulgated. It is also used to cure arterial hypertension; diabetes, renal diseases and cardiovascular diseases in different countries. Due to its medicinal paramountcy and use in food industry, perfume manufacturing, soaps and creams its cultivation should be elevated to meet its rising demand in the entire world.

## REFERENCES

- Alireza Yousofi, Saeed Daneshmandi, Neda Soleimani, Kambiz Bagheri and Mohammad Hossein Karimi (2012). Immunomodulatory effect of Parsley (*Petroselinum crispum*) essential oil on immune cells: Mitogenactivated splenocytes and peritoneal macrophages, Immunopharmacology and Immunotoxicology, Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. **34**(2): 303-308.
- Ambrus CM, Ambrus JL, Horvath C Pedersen, Sharma H, Kant SC, Mirand E, Guthrie R and Paul T (1978). Phenylalanine depletion for the management of phenylketonuria: Use of enzyme reactors with immobilized enzymes. *Science*, **201**: 837-839.
- Anand NK, Sharma ND and Gupta SR (1981). Coumarins from *Apium petroselinum* seeds. *Natl Acad Sci. Lett.*, **4**: 249-251.
- Azzini E, Bugianesi R, Romano F, Di Venere D, Miccadei S, Durazzo A, Foddai MS, Catasta G, Linsalata V and Maiani G (2007). Absorption and metabolism of bioactive molecules after oral consumption of cooked edible heads of *Cynara scolymus* L (cultivar Violetto di Provenza) in human subjects: A pilot study. *Br. J. Nutr.*, **97**: 963-969.
- Baytop T (1984). Therapy with medicinal plants in Turkey (Past and Present), Istanbul University Yayinlari, Turkey, p.3255

- Birt DF, Mitchell D, Gold B, Pour P and Pinch HC (1997). Inhibition of ultraviolet light-induced skin carcinogenesis in SKH-1 mice by apigenin, a plant flavonoid. *Anticancer Res.*, **17**: 85-91.
- Bolkent S, Yanardag R, Ozsoy-Sacan O and Karabulut-Bulan O (2004). Effects of Parsley (*Petroselinum crispum*) on the liver of diabetic rats. A morphological and biochemical study. *Phytother. Res.*, **18**: 996-999.
- Branković S, Kitić D, Radenković M, Veljković S, Milica Kostić, Miladinović B and Pavlović D (2010). Hypotensive and cardioinhibitory effects of the aqueous and ethanol extracts of celery (*apium graveolens*, apiaceae) *Acta. Medica Medianae*, **49**(1): 13-16.
- Campanella L, Bonanni A, Favero G and Tomassetti M (2003). Determination of antioxidant properties of aromatic herbs, olives and fresh fruit using an enzymatic sensor. *Anal. Bioanal. Chem.*, **375**: 1011-1016.
- Charly Masanetz and Werner Grosch (1998). Key Odorants of Parsley Leaves (*Petroselinum crispum* [Mill.] Nym. ssp. *crispum*) by Odour-Activity Values. *Flav and Frag J*, **13**: 115-124.
- Chaves DS, Frattani FS, Assafim M, de Almeida AP, de Zingali RB, Costa SS and Chaves (2011). Phenolic chemical composition of *Petroselinum crispum* extract and its effect on haemostasis. *Nat Prod. Commun.*, **6**(7): 961-964.
- Christoph Kirsch, Elke Logemann, Bernadette Lippok, Elmon Schmelzer and Klaus Hahlbrock (2001). A highly specific pathogen-responsive promoter element from the immediate-early activated CMPG1 gene in *Petroselinum crispum*. *The Plant Journal*, **26**(2): 217-227.
- Dahiru D, Sini JM and John-Africa L (2006). Antidiarrhoeal activity of *Ziziphus mauritiana* root extract in rodents. *Afr. J. Biotechnol.*, **5**(10): 941-945.
- Damien Dorman HJ, Tiina A Lantto, Atso Raasmaja and Raimo Hiltunen (2011). Antioxidant, pro-oxidant and cytotoxic properties of parsley. *Food Funct.*, **2**: 328-337.
- Darias V, Martin-Herrera D and Abdala S (2001). Plants used in urinary pathologies in the Canary islands. *Pharm Biol*, **39**: 170.
- Davey MW, Bauw G and Montagu MV (1996). Analysis of ascorbate in plant tissue by high performance capillary zone electrophoresis. *Anal Biochem.*, **239**: 8-19.
- Díaz-Maroto MC, Pérez-Coello MS and Cabezudon MD (2002). Effect of different drying methods on the volatile components of parsley (*Petroselinum crispum* L). *Eur. Food Res. Technol.*, **215**: 227-230.
- Dirk Nennstiel, Dierk Scheel and Thorsten Nürnberger (1998). Characterization and partial purification of an oligopeptide elicitor receptor from parsley (*Petroselinum crispum*), *FEBS Letters*, **431**(3): 405-410.
- Dounia Gadia, Mohamed Bnouhama, Mohammed Aziza, Abderrahim Ziyayata, Abdelkhaleq Legssyera, Chantal Legrand, Françoise Fauvel Lafeveb and Hassane Mekhfi (2009). Parsley extract inhibits in vitro and ex vivo platelet aggregation and prolongs bleeding time in rats. *J Ethnopharmacol*, **125**(1): 170-174.
- Eddouks M, Maghrani M, Lemhadri A and Ouahidi MLH (2002). Jouad Ethnopharmacological survey of medicinal plants used for the treatment of diabetes mellitus, hypertension and cardiac diseases in the south-east region of Morocco (Tafilalet). *J Ethnopharmacol*, **82**: 97-103.
- Engelmann C, Blot E, Panis Y, Bauer S, Trochon V, Nagy HJ, Lu H and Soria C (2002). Apigenin-strong cytostatic and antiangiogenic action *in vitro* contrasted by lack of efficacy *in vivo*. *Phytomedicine*, **9**: 489-495.
- Fahad A Alyami and Danny M Rabah (2011). Effect of drinking parsley leaf tea on urinary composition and urinary stones' risk factors. *Saudi J Kidney Dis Transpl.*, **22**(3): 511-514.
- Fatemeh Haidari, Seid AK, Majid Mammad Shahi, Soltan-Ali Mahboob and Mohammad-Reza Rashidi (2011). Effects of Parsley (*Petroselinum crispum*) and its Flavonol Constituents, kaempferol and quercetin, on serum uric acid levels, biomarkers of oxidative stress and liver xanthine oxidoreductase activity in oxonate-induced hyperuricemic rats. *Iran J. Pharm. Res.*, **10**(4): 811-819.
- Fejes S, Blazovics A, Lemberkovics E, Petri G, Szoke E and Kery A (2000). Free radical scavenging and membrane protective effects of methanol extracted fractions of parsley. *Acta-Alimentaria.*, **29**: 81-87.
- Fernandez de Simon B, Perez Ilzarbe J, Hernandez T, Gomez Cordoves C and Estrella I (1992). Importance of phenolic compounds for the characterization of fruit juices. *J. Agri Food Chem*, **40**: 1531-1535.
- Fraga CG, Martino VS, Ferraro GE, Coussio JD and Boveris A (1987). Flavonoids as antioxidants evaluated by *in vitro* and *in situ* liver chemiluminescence. *Bioch Pharmacol*, **36**: 717-720.
- Folling I (1994). The discovery of phenylketonuria. *Acta. Paediatr (Suppl.)*, **407**: 4-10.
- Gregory LH, Ken MR and Steven JS (2012). Endogenous enzymes, heat and pH affect flavone profiles in Parsley (*Petroselinum crispum* var *neapolitanum*) and Celery (*Apium graveolens*) during juice processing. *J. Agr. Food Chem.*, **60**: 202-208.
- Gupta SS (1994). Prospects and perspectives of natural plant products in medicine. *Indian J. Pharmacol.*, **26**(1): 1-12.
- Hahlbrock K and Scheel D (1989). Annual review of plant physiology and plant molecular biology. *Annu. Rev. Plant Biol.*, **40**: 347.
- Hempel J, Pforte H, Raab B, Engst W, Bohm H and Jacobasch G (1999). Flavonols and flavones of parsley cell suspension culture change the antioxidative capacity of plasma in rats. *Nahrung*, **43**: 201-204.
- Holger R and Georg ES. (2004). Structural basis for the

- entrance into the phenylpropanoid metabolism catalyzed by phenylalanine ammonia-lyase. *Plant Cell*, **16**(12): 3426-3436.
- Hoskins JA, Holiday SB and Greenway AM (1984). The metabolism of cinnamic acid by healthy and phenylketonuric adults: A kinetic study. *Biomed. Mass Spectrom.*, **11**: 296-300.
- Hui Zhang, Feng Chen, Xi Wang and Hui-Yuan Yao (2006). Evaluation of antioxidant activity of parsley (*Petroselinum crispum*) essential oil and identification of its antioxidant constituents. *Food Res. Int.*, **39**: 833-839.
- Ikeda K, Schiltz E, Fujii T, Takahashi M, Mitsui K, Kodera Y, Matsushima A, Inada Y, Schulz GE and Nishimura H (2005). Phenylalanine ammonia-lyase modified with polyethylene glycol: Potential therapeutic agent for phenylketonuria. *Amino Acids*. **29**(3): 283-287.
- Irzaie NM, Amabi D, Moazedi AA and Seyyednejad SM (2010). The role of and adrenergic receptors in the spasmolytic effects on rat ileum of *Petroselinum crispum latifolium* (Parsley). *Asian Pac. J. Trop. Med.*, **3**(11): 866-870.
- Jervis DA (1960). Detection of heterozygotes for phenylketonuria. *Clin. Chim. Acta.*, **5**: 471-476.
- Janssen K, Mensink RP, Cox FJ, Harryvan JL, Hovenier R, Hollman PC and Katan MB (1998). Effects of the flavonoids quercetin and apigenin on hemostasis in healthy volunteers: results from an *in vitro* and a dietary supplement study. *Am. J. Clin. Nutr.*, **67**: 255-262.
- Jens H, Olaf B, JuÈrgen S, Victor W, Klaus H and Dieter S (1999). Accumulation of phthalides in elicitor-treated cell suspension cultures of *Petroselinum crispum*. *Phytochemistry*, **51**: 629-635.
- Jeong YJ, Choi YJ, Choi JS, Kwon HM, Kang SW, Bae JY, Lee SS, Kang JS, Han SJ and Kang YH (2007). Attenuation of monocyte adhesion and oxidised LDL uptake in luteolin-treated human endothelial cells exposed to oxidised LDL. *Br. J. Nurt.*, **97**: 447-457.
- Jimenez-Alvarez D, Giuffrida F, Golay PA, Cotting C, Lardeau A and Keely BJ (2008). Antioxidant activity of oregano, parsley and olive mill wastewaters in bulk oils and oil in water emulsions enriched in fish oil. *J. Agric. Food Chem.*, **56**(16): 7151-7159.
- Jouad H, Haloui M, Rhiouani H, Hilaly J El and Eddouks M (2001). Ethnobotanical survey of medicinal plants used for the treatment of diabetes, cardiac and renal diseases in the North centre region of Morocco (Fez-Boulemane). *J. of Ethnopharmacol*, **77**: 175-182.
- Jung HP, Sen A and Grosch W (1992). Evaluation of potent ordants in Parsley Leaves by aroma Extract Dilution Analysis, *Lebensm.-Wiss. -Technol.* **25**: 55-60.
- Justesen U, Knuthsen P and Leth T (1998). Quantitative analysis of flavonols, flavones, and flavanones in fruits, vegetables and beverages by HPLC with photodiode array and mass spectrometric detection. *Journal of Chromatography*, **799**: 101-110.
- Kasting R, Anderson J and von Sydow E (1972). Volatile constituents in leaves of parsley. *Phytochem*, **11**: 2277-2282.
- Keser G and Buyuk G (2012). Effects of wastewater irrigation on chemical and physical properties of *Petroselinum crispum*. *Biol Trace Elem Res.*, **146**(3): 369-375.
- Kitagawa I, Yoshikawa M and Yoshioka I (1976). sapogenin and sapogenol XIII. Structures of three soyabean saponins: soyssaponin II, & soyasaponin III. *Chem. Pharm. Bull.*, **24**: 121-129.
- Kreydiyyeh SI and Usta J (2002) Diuretic effect and mechanism of action of parsley, *J. Ethnopharmacol.*, **79**: 353-7.
- Kreydiyyeh SI Usta J, Kaouk I and Al-Sadi R (2001). The mechanism underlying the laxative properties of parsley extract. *J. Phytomed.*, **8**(5): 382-8.
- Kühnau J (1976). The flavonoids, A class of semi-essential food components: their role in human nutrition. *World Rev Nut Diet*, **24**: 117-191.
- Kuo-Hsiung L and Soine TOJ (1969). Coumarins X: Spectral studies on some linear furanocoumarins . *Pharm. Sci.*, **58**: 681-683.
- Kuo ML, Lee KC and Lin JK (1992). Genotoxicities of nitropyrenes and their modulation by apigenin, tannic acid, ellagic acid and indole-3-carbinol in the Salmonella and CHO systems. *Mutation Research*, **270**: 87-95.
- Lechtenberg M, Zumdick S, Gerhards C, Schmidt TJ and Hensel A (2007). Evaluation of analytical markers characterising different drying methods of parsley leaves (*Petroselinum crispum* L). *Pharmazie*, **62**(12): 949-54.
- Lee SJ, Son KH, Chang HW, Do JC, Jung KY, Kang SS and Kim HP (1993). Antiinflammatory activity of naturally occurring flavone and flavonol glycosides, *Archives of Pharmacol Research*, **16**: 25-28
- Lopez MG, Sancheze-Medoza IR and Ochoa-Alejo N (1999). Comparative study of volatile components and fatty acids of plants and *in vitro* cultures of parsley (*Petroselinum crispum*) (Mill nym ex hill). *J. Agric Food Chem.*, **47**: 3292-3296.
- Mabile L, Bruckdorfer KR and Rice-Evans C (1999). Moderate supplementation with natural alpha-tocopherol decreases platelet aggregation and low-density lipoprotein oxidation. *Atherosclerosis*, **147**: 177-185.
- MacLeod AJ, Snyder CH and Subramanian G (1985). Volatile aroma constituents of parsley leaves. *Phytochem.*, **24**: 2623-2627.
- Manderfeld MM, Schafer HW, Davidson PM and Zottola EA (1997). Isolation and identification of antimicrobial furanocoumarins from parsley. *J. Food Prot.*, **60**: 72-77.
- Manerfeld MM, Schafer HW, Davidson PM and Zottola EA (1997). Isolation and identification of antimicrobial

- furocoumarins from parsley. *J. food Prot.*, **60**: 72-77.
- Manthey JA and Guthrie N (2002). Antiproliferative activities of citrus flavonoids against six human cancer cell lines. *J. Agric. Food Chem.*, **50**: 5837-5843.
- Marczal G, Balogh M and Verzar-Petri G (1997). Phenol-ether components of diuretic effect in Parsley I. *Acta. Agron Acad. Sci. Hung.*, **26**: 7-13.
- Markham KR, Ternai B, Stanley R, Geiger H and Mabry TL (1978). <sup>13</sup>C NMR studies of flavonoids 3 naturally occurring flavonoid glycosides and their acylated derivatives. *Tetrahedron*, **34**: 1389-1397.
- Masanetz C and Grosch W (1998). Key odorants of parsley leaves (*Petroselinum crispum* [Mill.] Nym. ssp. *crispum*) by odouractivity values. *Flavour Fragr J.*, **13**: 115-124.
- Masayuki Y, Hisashi M, Toshiaki U, Hiroshi S, Akinobu K and Yuzo K (2000). Medicinal foodstuffs. Phytoestrogens from the aerial part of *Petroselinum crispum* mill (parsley) and Structures of 6-Acetylapiin and a New Monoterpene Glycoside Petroside, *Chem. Pharm. Bull.*, **48**(7): 1039-1044.
- Massey LK, Liebman M, Kynast-Gales SA (2005). Ascorbate Increases Human Oxaluria and Kidney Stone Risk, *J. Nutr.*, **135**: 1673-1677.
- Matern U (1991). Coumarins and other phenylpropanoid compounds in the defense response of plant cells. *Planta Medica*, **57**: 15.
- Mekhfi H, Haouari M El , Legssyer A, Bnouham M, Aziz M, Atmani F, Remmal A and Ziyat A (2004). Platelet anti-aggregant property of some Moroccan medicinal plants. *J Ethnopharmacol*, **94**: 317-322.
- Moazedi AA, Mirzaie-Damabi N, Seyyednejad SM, Zadkarami MR and Amirzargar A (2007). Spasmolytic effect of *Petroselinum crispum* (Parsley) on rats ileum at different calcium chloride concentrations. *Pak. J. Biol. Sci.*, **10**(22): 4036-4042.
- Newall C, Anderson L and Phillipson J (1996). Herbal Medicine: A Guide for Health-Care Professionals. Pharmaceutical Press, London, pp.203-204
- Nicholas C, Batra S, Vargo MA, Voss OH, Gavrillin MA, Wewers MD, Guttridge DC, Grotewold E and Doseff AI (2007). Apigenin blocks lipopolysaccharide-induced lethality in vivo and proinflammatory cytokines expression by inactivating NF-κB through the suppression of P65 phosphorylation. *J. Immunol.*, **179**: 7121.
- Nielsen SE, Young JF, B Daneshvar, ST Lauridsen, P Knuthsen and B Sandstrom (1999). Effect of parsley intake on urinary apigenin excretion, blood antioxidant enzymes and biomarkers for oxidative stress in human subjects. *Br. J. Nutr.*, **81**: 447-455.
- Noel PH, Pugh JA and Larme AC (1997). The use of traditional plant medicines for non insulin dependent diabetes mellitus in South Texas. *Phytotherapy Research*, **11**: 512-517.
- Okuyama T, Hosoyama K, Hiraga Y, Kurono G and Takemoto T (1978). The constituents of *Osmunda* spp. II: A new flavonol glycoside of *Osmunda asiatica*. *Chem. Pharm. Bull.*, **26**: 3071-3074.
- Ozsoy-Sacan O, Yanardag R, Orak H, Ozgey Y, Yarat A and Tunali T (2006). Effects of parsley (*Petroselinum crispum*) extract versus glibornuride on the liver of streptozotocin-induced diabetic rats. *J. Ethnopharmacol.*, **104**(1-2): 175-181.
- Ozturk Y, CHK Baser and S Aydin (1991). Hepatoprotective (antihepatotoxic) plants in Turkey. Proceedings of the 9<sup>th</sup> Symposium on Plant Drugs. Eskisehir Turkey, pp.40-50.
- Pace-Asciak CR, Hahn S, Diamandis EP, Soleas G, Goldberg DM (1995). The red wine phenolics trans-resveratrol and quercetin block human platelet aggregation and eicosanoid synthesis: Implications for protection against coronary heart disease. *Clinica Chimica Acta.*, **235**: 207-219.
- Pearson DA, Paglieroni TG, Rein D, Wun T, Schramm DD, Wang JF, Holt RR, Gosselin R, Schmitz HH and Keen CL (2002). The effect of flavonol-rich cocoa and aspirin on ex vivo platelet function. *Thrombosis Research*, **106**: 191-197.
- Peter YY Wong and David D Kitts (2006). Studies on the dual antioxidant and antibacterial properties of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) extracts. *Food Chem*, **97**: 505-515.
- Petropoulos SA, Daferera D, Akoumianakis CA, Passam HC and Polissiou MG (2004). The effect of sowing date and growth stage on the essential oil composition of three types of parsley *Petroselinum crispum*. *J. Sci Food Agr*, **84**: 1606-1610.
- Pino JA, Rosada A and Fuentes V (1997). Herb oil of parsley (*Petroselinum crispum* Mill.) from Cuba. *J. Essent Oil Res.*, **9**: 241-242.
- Redaelli C, Formentini L and Santaniello E (1980). Apigenin 7-glucoside and its 2- and 6-acetates from ligulate flowers of *Matricaria chamomilla*. *Phytochemistry*, **19**: 985-986.
- Rein D, Paglieroni TG, Pearson DA, Wun T, Schmitz HH, Gosselin R and Keen CL (2000). Cocoa and wine polyphenols modulate platelet activation and function. *Nutrition*, **130**: 2120-2126.
- Sakakibara H, Honda Y, Nakagawa S, Ashida H and Kanazawa K (2003). Simultaneous determination of all polyphenols in vegetables, fruits, and teas. *J. Agric. Food Chem.*, **51**: 571-581.
- Scheel D, Hau KD, Jahnen W and Hahlbrock K (1986). In B. Lugtenberg, Recognition in microbe-plant symbiotic and pathogenic interactions. Springer-Verlag, Berlin, p.325.
- Sebastian Bartsch and Uwe T Bornscheuer (2010). Mutational analysis of phenylalanine ammonia lyase to improve reactions rates for various substrates. *Proteins Eng Des SEL*:**23**(12): 929-933.
- Simon JE and Quinn J (1988). Characterization of essential oil of parsley. *J. Agric Food Chem.*, **36**: 467-47.

- St. Louis (1991). Review of Natural Products. MO: Facts and Comparisons.
- Subramoniam P and Pushpangadan P (1999). Development of phytomedicines for liver disease. *Indian J. Pharmacol.*, **31**(3): 166-175.
- Nürnberg, Nennstiel D, Jabs T, Sacks WR, Hahlbrock K and Scheel D (1994). High-affinity binding of a fungal oligopeptide at parsley plasma membranes triggers multiple defense responses. *Cell*, **78**: 449-460.
- Tahan M and Bayram I (2011). Effect of using black cumin (*Nigella sativa*) and parsley (*Petroselinum crispum*) in laying quail diets on egg yield, egg quality and hatchability. *Archiva Zootechnica*, **14**(4): 39-44.
- Tunali T, Yarat A, Yanardag R, Ozcelik F, Ozsoy O, Ergenekon G and Emekli N (1999). Effect of parsley (*Petroselinum crispum*) on the skin of STZ induced diabetic rats. *Phyto Res.*, **13**: 138-141.
- Tyler V (1994). Herbs of Choice. Pharmaceutical Product Press, New York, pp.75-76.
- Ueda, H, Yamazaki C and Yamazaki M (2004). A hydroxyl group of flavonoids affects oral anti-inflammatory activity and inhibition of systemic tumor necrosis factor- $\alpha$  production. *Biosci. Biotechnol. Biochem.*, **68**: 119-125.
- Van Acker SABE, van den Berg DJ, Tromp MNJL, Griffioen DH, van Bennekom WP, van der Vijgh WJF and Bast A (1996). Structural aspects of antioxidant activity of flavonoids. *Free Radic. Bio. Med.*, **20**: 331-342.
- Vargo MA, Voss OH, Poustka F, Cardounel AJ, Grotewold E and Doseff AI (2006). Apigenin-induced-apoptosis is mediated by the activation of PKC  $\delta$  and caspases in leukemia cells. *Biochem. Pharmacol.*, **72**: 681-692.
- Vernon F and Richard HMJ (1983). *Lebensm Wiss u Technol.*, **16**: 32-35.
- Vokk R, Lõugas T, Mets K and Kravets M (2011). Dill (*Anethum graveolens* L.) and Parsley (*Petroselinum crispum* (Mill) Fuss) from Estonia: Seasonal differences in essential oil composition. *Agron Res*, **9**(2): 515-520.
- Vora Shreya R, Patil Rahul B and Pillai Meena M (2009). Protective effects of *Petroselinum crispum* (Mill) Nyman ex A.W Hill leaf extract on D-galactose-induced oxidative stress in mouse brain. *Indian J. Exp Biol*, **47**(05): 338-342.
- Wahba NM, Ahmed AS and Ebraheim ZZ (2010). Antimicrobial effects of pepper, parsley and dill and their roles in the microbiological quality enhancement of traditional Egyptian Kareish cheese. *Foodborne Pathog Dis. Apr.*, **7**(4): 411-418.
- Walle T, Kawamori Ta N, Wen T, Tsuji X and Walle PA (2007). Cancer chemopreventive properties of orally bioavailable flavonoids-methylated versus unmethylated flavones. *Biochem. Pharmacol.*, **73**: 1288-1296.
- Wei A and Shibamoto T (2007). Antioxidant activities and volatile constituents of various essential oils. *J. Agric Food Chem.*, **55**(5): 1737-1742.
- Wei H, Tye L, Bresnick E and Birt DF (1989). Inhibitory effect of apigenin, a plant flavonoid, on epidermal ornithine decarboxylase and skin tumor promotion in mice. *Cancer Research*, **50**: 499-502.
- Wong PYY and Kitts DD (2006). Studies on the dual antioxidant and antibacterial properties of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) extracts. *Food Chem.*, **97**: 505-515.
- Yakushijin K, Tohshima T, Suzuki R, Murata H, Lu ST and Furukawa H (1983). Studies on the constituents of the plants of Illicium species. II. Structures of phenolic components. *Chem. Pharm. Bull.*, **31**: 2879-2883.
- Yanardag R, Bolkent S, Tabakoğlu-Oğuz A and Ozsoy-Saçan O (2003). Effects of *Petroselinum crispum* extract on pancreatic B cells and blood glucose of streptozocin induced diabetic rats. *Biol Pharm Bull*, **26**: 1206-1207.
- Yvonne Helen Gebhardt, Simone Witte, Holger Steuber, Ulrich Matern and Stefan Martens (2007). Evolution of Flavone Synthase I from Parsley Flavanone 3 $\beta$ -Hydroxylase by Site-Directed Mutagenesis. *Plant Physiol*, **144**(3): 1442-1454.
- Ziyyat A, Legssyer A, Mekhfi H, Dassouli A, Serhrouchni M and Benjelloun W (1997). Phytotherapy of hypertension and diabetes in oriental Morocco. *J Ethnopharmacol*, **58**: 45-54.